

Solving Mathematical Programs with Equilibrium Constraints as Nonlinear Programming: A New Framework

Songqiang Qiu · Zhongwen Chen

Received: date / Accepted: date

Abstract We present a new framework for the solution of mathematical programs with equilibrium constraints (MPECs). In this algorithmic framework, an MPECs is viewed as a concentration of an unconstrained optimization which minimizes the complementarity measure and a nonlinear programming with general constraints. A strategy generalizing ideas of Byrd-Omojokun's trust region method is used to compute steps. By penalizing the tangential constraints into the objective function, we circumvent the problem of not satisfying MFCQ. A trust-funnel-like strategy is used to balance the improvements on feasibility and optimality. We show that, under MPEC-MFCQ, if the algorithm does not terminate in finite steps, then at least one accumulation point of the iterates sequence is an S-stationary point.

Keywords Mathematical programs with equilibrium constraints · nonlinear programs · trust-funnel-like methods · S-stationarity · Global convergence

Mathematics Subject Classification (2000) 90C30 · 90C55 · 65K05

1 Introduction

We consider the Mathematical programs with equilibrium constraints (MPECs) in the form of

$$\begin{aligned} \min \quad & f(x) \\ \text{s.t.} \quad & g(x) \geq 0, \quad h(x) = 0, \\ & G(x) \geq 0, \quad H(x) \geq 0, \\ & G(x)^T H(x) = 0, \end{aligned} \tag{1}$$

This work was supported by the Fundamental Research Funds for the Central Universities 2014QNA62

Songqiang Qiu, Corresponding author
School of Sciences, China University of Mining and Technology
Xuzhou, China
sqqiu@cumt.edu.cn

Zhongwen Chen
School of Mathematical Sciences, Soochow University
Suzhou, China
zwchen@suda.edu.cn

where $f : \mathbb{R}^n \rightarrow \mathbb{R}$, $g : \mathbb{R}^n \rightarrow \mathbb{R}^m$, $h : \mathbb{R}^n \rightarrow \mathbb{R}^p$, $G, H : \mathbb{R}^n \rightarrow \mathbb{R}^q$ are continuously differentiable. This problem is also called mathematical programs with complementarity constraints (MPCCs). Problems of this type can be used to model numerous economics or engineering applications [1, 2]. However, in a theoretical view, it has been shown that MPECs does not satisfy constraint qualifications [3, 4] at any feasible point. As a result, the linearized constraints may be infeasible even near a feasible point, which causes it problematic to solve MPECs using standard algorithm for nonlinear programmings.

Several methods have been proposed to handle such problems. A number of algorithms are characterized by embedding an MPEC within a one parameter family of nonlinear programs. This type of approaches includes smoothing algorithms [5, 6], the regularization or relaxation methods [7, 8] and penalty methods [9, 10, 8]. Some other methods use the idea of piecewise sequential quadratic programming, for example, the piecewise sequential quadratic programming (PSQP) method [11, 1, 12], the penalty decomposition methods of Ref. [13].

A quite different but interesting way for solving MPECs is to solve it as nonlinear programming (NLP). Several researchers have studied some methods of this type, both in theory and in numerical performance. SQP-based approaches have been presented in [14, 15, 16, 17] and etc while interior-point-based methods in [18, 19, 20] and etc. Numerical performance on MacMPEC [21] were reported in [18, 22, 19] and the results were encouraging.

In this paper, we present a new SQP-based algorithmic scheme for solving MPECs. The main ideas of our algorithm is to view MPECs (1) as the following two sequential problems

$$\min \quad \theta^C(x) = |G(x)^T H(x)| \quad (2)$$

and

$$\begin{aligned} \min \quad & f(x) \\ \text{s.t.} \quad & g(x) \geq 0, \quad h(x) = 0, \\ & G(x) \geq 0, H(x) \geq 0, \end{aligned} \quad (3)$$

Then the process of solving MPECs is decomposed into two steps of which the first step is to improve complementarity and the second step is to solve a quadratic programming model for the nonlinear programming (3). To trade off the progresses achieved by these two steps, a penalized Byrd-Omojukun-like idea is used.

We use a line search trust-funnel-like strategy to balance the improvements on feasibility and optimality. A maximal infeasibility measure is set and no iterate is allowed to violate the restriction of it. A line search is performed with certain acceptance criteria in accordance with the comparison between infeasibility measure and reduction on linearize objective function. We show that, under MPEC Mangasarian Fromovitz constraint qualification (MPEC-MFCQ), if the algorithm does not terminate finitely, then at least one accumulation point of the sequence of iterates is strong stationary.

The paper is structured as follows. In section 2, we introduce some background knowledges for the constraint qualifications and stationarity of MPECs. In section 3, we describe the nature of the algorithm in detail. In section 4, we present the global convergence analysis for the given algorithm. Finally, we give some comments on the further work.

Notations: We use $\|\cdot\|$ to denote the Euclidean norm $\|\cdot\|_2$. Subscript k refers to iteration indices and superscript (i)

2 Background knowledge

The active sets at a feasible point $\tilde{x}$ is defined as

$$\begin{aligned} I_g &\stackrel{\text{def}}{=} \{i = 1, \dots, m \mid g^{(i)}(\tilde{x}) = 0\}, \\ I_G &\stackrel{\text{def}}{=} \{i = 1, \dots, q \mid G^{(i)}(\tilde{x}) = 0\}, \\ I_H &\stackrel{\text{def}}{=} \{i = 1, \dots, q \mid H^{(i)}(\tilde{x}) = 0\}. \end{aligned}$$

Next we introduce some stationarity concepts for MPECs. The first definition of stationarity is the definition of B-stationarity.

Definition 2.1 A feasible point $\tilde{x}$ of (1) is called a Bouligand- or B-stationary point, if $d = 0$ solves the following linear programming with equilibrium constraints (LPECs)

$$\begin{aligned} \min \quad & \nabla f(\tilde{x})^T d \\ \text{s.t.} \quad & g(\tilde{x}) + \nabla g(\tilde{x})^T d \geq 0, \\ & h(\tilde{x}) + \nabla h(\tilde{x})^T d = 0, \\ & G(\tilde{x}) + \nabla G(\tilde{x})^T d \geq 0, \\ & H(\tilde{x}) + \nabla H(\tilde{x})^T d \geq 0, \\ & (G(\tilde{x}) + \nabla G(\tilde{x})^T d)^T (H(\tilde{x}) + \nabla H(\tilde{x})^T d) = 0. \end{aligned}$$

Checking B-stationarity is difficult in general because it involves the solution of an LPEC which is a combinatorial problem. However, B-stationarity is implied by the following S-stationarity, which is more restrictive but much easier to check.

Definition 2.2 A feasible point $\tilde{x}$ of (1) is called a Strong- or S-stationary point, if $d = 0$ solves the following linear programming

$$\begin{aligned} \min \quad & \nabla f(\tilde{x})^T d \\ \text{s.t.} \quad & g(\tilde{x}) + \nabla g(\tilde{x})^T d \geq 0, \\ & h(\tilde{x}) + \nabla h(\tilde{x})^T d = 0, \\ & \nabla G^{(i)}(\tilde{x})^T d = 0, \quad i \in I_G \setminus I_H, \\ & \nabla H^{(i)}(\tilde{x})^T d = 0, \quad i \in I_H \setminus I_G \\ & \nabla G^{(i)}(\tilde{x})^T d \geq 0, \quad \nabla H^{(i)}(\tilde{x})^T d \geq 0, \quad i \in I_G \cap I_H. \end{aligned} \tag{4}$$

Note that (4) is the linearized approximation of the following nonlinear programming which is referred to as the relaxed nonlinear programming (RNLP) for (1):

$$\begin{aligned} \min \quad & f(x) \\ \text{s.t.} \quad & g(x) \geq 0, \quad h(x) = 0, \\ & G^{(i)}(x) = 0, \quad i \in I_G \setminus I_H, \quad (RNLP) \\ & H^{(i)}(x) = 0, \quad i \in I_H \setminus I_G, \\ & G^{(i)}(x) \geq 0, \quad H^{(i)}(x) \geq 0, \quad i \in I_G \cap I_H. \end{aligned} \tag{5}$$

By introduction of the following Lagrangian function [7]

$$\mathcal{L}(x_k, \lambda_k, \mu_k, \nu_k, \xi_k) = f_k - \lambda_k^T g_k + \mu_k^T h_k - \nu_k^T G_k - \xi_k^T H_k,$$

we can give the definition of weaker stationarity and the uniform definitions of C-stationarity, M-stationarity and S-stationarity.

Definition 2.3 [23] A feasible point $\tilde{x}$ of MPEC (1) is called weakly stationary if there exists MPEC multipliers $\tilde{\lambda} \geq 0$, μ , $\tilde{\nu}$ and $\tilde{\xi}$ satisfying

$$\nabla_x \mathcal{L}(\tilde{x}, \tilde{\lambda}, \tilde{\mu}, \tilde{\nu}, \tilde{\xi}) = 0, \quad \tilde{\lambda}^T g(\tilde{x}) = 0, \quad \tilde{\nu}^T G(\tilde{x}) = 0, \quad \tilde{\xi}^T H(\tilde{x}) = 0. \quad (6)$$

And $\tilde{x}$ is called

- a C-stationarity if $\tilde{\nu}^{(i)} \tilde{\xi}^{(i)} \geq 0$ for all $i \in I_G(\tilde{x}) \cap I_H(\tilde{x})$;
- an M-stationarity point if for all $i \in I_G(\tilde{x}) \cap I_H(\tilde{x})$, either $\tilde{\nu}^{(i)} \tilde{\xi}^{(i)} > 0$ or $\tilde{\nu}^{(i)} \tilde{\xi}^{(i)} = 0$;
- an S-stationarity point if

$$\tilde{\nu}^{(i)} \tilde{\xi}^{(i)} \geq 0 \text{ for all } i \in I_G(\tilde{x}) \cap I_H(\tilde{x}). \quad (7)$$

Clearly, the definition of S-stationarity here is equivalent to definition 2.2. S-stationarity and B-stationarity are equivalent in the case that the following MPEC-LICQ is satisfied.

Definition 2.4 The MPEC-LICQ is satisfied at the point $\tilde{x}$ if the following set of vectors is linearly independent:

$$\{\nabla g^{(i)}(\tilde{x}) | i \in I_g\} \cup \{\nabla h^{(i)}(\tilde{x}) | i = 1, 2, \dots, p\} \cup \{\nabla G^{(i)}(\tilde{x}) | i \in I_G\} \cup \{\nabla H^{(i)}(\tilde{x}) | i \in I_H\}.$$

In other words, the linear independence constraint qualification (LICQ) is satisfied for RNLP (5).

When the MPEC-LICQ condition holds, we have the following first-order necessary conditions, see, for instance, [1] and [24, Thm. 2].

Theorem 2.1 Suppose that $\tilde{x}$ is a local minimizer of (1) and that the MPEC-LICQ condition holds at $\tilde{x}$. Then $\tilde{x}$ is strongly stationary; and the multiplier vector $(\tilde{\lambda}, \tilde{\mu}, \tilde{\nu}, \tilde{\xi})$ that satisfies (6) and (7) is unique.

Our global convergence analysis uses the following weaker Mangasarian-Fromovitz constraints qualification

Definition 2.5 The MPEC-MFCQ is satisfied at a feasible point $\tilde{x}$ of (1) if the MFCQ is satisfied for the RNLP (5); that is, if there exists a nonzero vector $d \in \mathbb{R}^n$ such that

$$\nabla G^{(i)}(\tilde{x})^T d = 0, \quad i \in I_G \setminus I_H, \quad \nabla H^{(i)}(\tilde{x})^T d = 0, \quad i \in I_H \setminus I_G, \quad (8)$$

$$\nabla h^{(i)}(\tilde{x})^T d = 0, \quad i = 1, 2, \dots, p, \quad (9)$$

$$\nabla g^{(i)}(\tilde{x})^T d > 0, \quad i \in I_g, \quad (10)$$

$$\nabla G^{(i)}(\tilde{x})^T d > 0 \text{ and } \nabla H^{(i)}(\tilde{x})^T d > 0, \quad i \in I_G \cap I_H; \text{ and} \quad (11)$$

$$\nabla G^{(i)}(\tilde{x}), \quad i \in I_G \setminus I_H, \quad \nabla H^{(i)}(\tilde{x}), \quad i \in I_H \setminus I_G, \quad \nabla h^{(i)}(\tilde{x}), \quad i = 1, 2, \dots, p \text{ are all linearly independent.} \quad (12)$$

3 Algorithm

3.1 Subproblems and steps

For convenience, we define $Q(x) = G(x)^T H(x)$. As we have mentioned in Section 1, the motivation to design this algorithm is the observation of decomposing the original problem (1) into problems (2) and (3). Correspondingly, the trial step is decomposed into a complementarity step, which aims to

improve the complementarity, and a nonlinear programming step, which looks for some improvement on (3). A complementary step is a step along the direction

$$s_k = -(\nabla Q_k^T)^\dagger Q_k = -\nabla Q_k (\nabla Q_k^T \nabla Q_k)^{-1} Q_k, \quad (13)$$

which is a least square solution of the linearized complementarity measure, i.e., s_k solves

$$\min |Q_k + \nabla Q_k s|^2.$$

A nonlinear programming step tries to make some improvement on (3). Following classic ideas of sequential quadratic programming (SQP) methods, such improvement can be obtained by solving

$$\begin{aligned} \min \quad & \nabla f_k^T(s_k + t) + \frac{1}{2}(s_k + t)^T B_k(s_k + t) \\ \text{s.t.} \quad & g_k + \nabla g_k^T(s_k + t) \geq 0, \\ & h_k + \nabla h_k^T(s_k + t) = 0, \\ & G_k + \nabla G_k^T(s_k + t) \geq 0, \\ & H_k + \nabla H_k^T(s_k + t) \geq 0. \end{aligned} \quad (14)$$

A side-effect of the solution t_k is that it could jeopardize the infeasibility reduction we have just obtained from step s_k . To weaken this negative effect, using Byrd-Omojokun's idea [25, 26], it makes good sense to replace (14) by the following quadratic programming:

$$\begin{aligned} \min \quad & \nabla f_k^T(s_k + t) + \frac{1}{2}(s_k + t)^T B_k(s_k + t) \\ \text{s.t.} \quad & g_k + \nabla g_k^T(s_k + t) \geq 0, \\ & h_k + \nabla h_k^T(s_k + t) = 0, \\ & G_k + \nabla G_k^T(s_k + t) \geq 0, \\ & H_k + \nabla H_k^T(s_k + t) \geq 0, \\ & \nabla Q_k^T t = 0. \end{aligned} \quad (15)$$

Unfortunately, the gradients of active constraints of this problem can not be linearly independent and no corresponding linear independency exists under MPEC-MFCQ. As a result, problem (15) may be infeasible even near a solution. This is mainly caused by the tangential constraint $\nabla Q_k^T t = 0$. To circumvent this problem, we penalize the constraint $\nabla Q_k^T t = 0$ by means of adding an item $\frac{1}{2u} t^T \nabla Q_k \nabla Q_k^T t$ to the objective function. Consequently, we get

$$\begin{aligned} \min \quad & \nabla f_k^T(s_k + t) + \frac{1}{2}(s_k + t)^T B_k(s_k + t) + \frac{1}{2u} t^T \nabla Q_k \nabla Q_k^T t \\ \text{s.t.} \quad & g_k + \nabla g_k^T(s_k + t) \geq 0, \\ & h_k + \nabla h_k^T(s_k + t) = 0, \\ & G_k + \nabla G_k^T(s_k + t) \geq 0, \\ & H_k + \nabla H_k^T(s_k + t) \geq 0, \end{aligned} \quad (16)$$

with $u > 0$. After simple calculations, we simplify (16) as

$$\begin{aligned} \min \quad & (\nabla f_k + B_k s_k)^T t + \frac{1}{2} t^T (B_k + \frac{1}{u} \nabla Q_k \nabla Q_k^T) t \\ \text{s.t.} \quad & g_k + \nabla g_k^T(s_k + t) \geq 0, \\ & h_k + \nabla h_k^T(s_k + t) = 0, \\ & G_k + \nabla G_k^T(s_k + t) \geq 0, \\ & H_k + \nabla H_k^T(s_k + t) \geq 0. \end{aligned} \quad (17)$$

In general, we have, by the classical convergence theorem of the quadratic penalty method [27]

Theorem 3.1 *Suppose that (15) is feasible at x_k , and that each $t_k(u_j)$ is the exact global minimizer of (17) defined above, and that $u_j \searrow 0$. Then every limit point $\tilde{t}_k$ of the sequence $\{t_k(u_j)\}$ is a global solution of the problem (15).*

However, the feasibility of (15) is not guaranteed even near a feasible point. Hence, problem (17) does not necessary lead to $\nabla Q_k^T t_k = 0$ for small u even in the limit case. Fortunately, we have observed that it is sufficient for the solution $t_k(u)$ of (17) to be close enough to the null space of ∇Q_k^T , i.e., to satisfy

$$|\nabla Q_k^T t_k(u)| \leq \delta_k, \quad (18)$$

where δ_k is a positive scalar which we will specify later. Generally speaking, the existence of u_k satisfying (18) is also not guaranteed if the sets of Lagrangian multipliers is unbounded. And, if the required u_k is very small, the algorithm may have to reduce u and solve the corresponding subproblem (17) many times, which is costly. We use a simple strategy to deal with this situation. If $t_k(u)$ fails to satisfy (18) while u becomes too small, say, $u < u_k^{\min}$ with $u_k^{\min}$ a small positive scalar, we stop reducing u , let $u_k := u$, $t_k = t_k(u_k)$ and scale t_k by $\gamma_k = \frac{\delta_k}{\|\nabla Q_k^T t_k\|}$. Here, we define $u_k^{\min}$ as

$$u_k^{\min} = \min\{\hat{u}, \kappa_u \delta_k^{\sigma_1}\},$$

where $\hat{u}, \kappa_u \in (0, 1)$ and $\sigma_1 > 1$. This correction “drags” t_k closer to the null space of ∇Q_k^T . Clearly, the scaled direction $\gamma_k t_k$ satisfies (18).

3.2 A trust-funnel-like strategy

Consider iteration k , starting from the iterate x_k . We use a trust-funnel-like strategy to balance the improvement of feasibility and optimality, and hence, to force the sequence of iterates converging to a first order stationary point. Trust funnel method was introduced by Gould and Toint [28]. Similar ideas, some of which follows directly the ideas of Gould and Toint, can be found in the methods proposed in [29, 30, 31, 32, 33, 34]. The key point of this kind of approaches is the control of infeasibility. A non-increasing limit $\theta_k^{\max}$ on infeasibility is introduced and no violation to this limit is allowed, i.e., there is $\theta_k \leq \theta_k^{\max}$ for all iterations. The infeasibility measure for the problem (1) is defined as

$$\theta_k := \theta_k^F + \theta_k^C,$$

where

$$\theta_k^F := \|(g_k)^-\| + \|h_k\| + \|(G_k)^-\| + \|(H_k)^-\|$$

and

$$\theta_k^C := |Q_k| = |G_k^T H_k|,$$

are used to qualify the dissatisfaction of the general constraints and complementarity constraint respectively. First, we compute a complementary step s_k by (13). Then, we solve (17) to obtain $t_k(u)$ with a given parameter u . If either $t_k(u)$ satisfies

$$-\nabla f(x_k)^T (s_k + t_k(u)) \geq \kappa_\theta \theta_k, \quad (19)$$

$$|\nabla Q_k^T t_k(u)| \leq \kappa_1 \min\{\theta_k^{\max}, \kappa_2\}, \quad (20)$$

with $\kappa_1 \in (0, 1/5)$, $\kappa_2 > 0$, or it satisfies

$$-\nabla f(x_k)^T(s_k + t_k(u)) < \kappa_\theta \theta_k, \quad (21)$$

$$|\nabla Q_k^T t_k(u)| \leq \kappa_3 \min\{\theta_k^{\sigma_2-1}, \kappa_2\} \theta_k, \quad (22)$$

with $\kappa_3 \in (0, 1/5)$ and $1 < \sigma_2 < \sigma_1$, then set $u_k = u$ and $t_k = t_k(u_k)$. Otherwise, reduce u and recompute $t_k(u)$. The choice of the parameter δ_k in (18) is clear now. We define

$$\delta_k = \begin{cases} \kappa_1 \min\{\theta_k^{\max}, \kappa_2\}, & \text{if (19) holds;} \\ \kappa_3 \min\{\theta_k^{\sigma_2-1}, \kappa_2\} \theta_k, & \text{if (21) holds.} \end{cases} \quad (23)$$

The whole search direction is thus given by $d_k = s_k + \gamma_k t_k$, where γ_k is defined as

$$\gamma_k = \begin{cases} \frac{\delta_k}{|\nabla Q_k^T t_k|}, & \text{if } u < u_k^{\min} \text{ and neither of (19) and (21) is satisfied,} \\ 1, & \text{otherwise.} \end{cases} \quad (24)$$

The condition (19) indicates that sufficient reduction on the linearized objective function is obtained along d_k . Therefore, it is reasonable to expect a satisfactory reduction on $f(x)$. The algorithm should put emphasis on reducing the value of objective function. Hence, we use a classical Amijio line search to find a step length α_k satisfying

$$f(x_k) - f(x_k + \alpha d_k) \geq -\alpha \rho \nabla f(x_k)^T d_k, \quad (25)$$

where $\rho \in (0, \min\{1 - 5\kappa_1, 1 - 5\kappa_3\})$ is fixed. The requirement on feasibility is rough in this case. We require

$$\theta(x_k + \alpha d_k) \leq \theta_k^{\max}. \quad (26)$$

In fact, it is based on the consideration of (26) that we require $t_k(u)$ to satisfy (20). In this case, we call d_k an f -step, x_k an f -iterate and k an f -iteration. If condition (21) holds, then reduction on objective function is not expectable, which is always caused by poor feasibility. Hence, the algorithm focuses on improving feasibility. Hereby, $\nabla Q_k^T t_k(u)$ is asked to admit (22). An Amijio line search is performed to find step length α_k satisfying

$$\theta(x_k + \alpha d_k) \leq (1 - \alpha \rho) \theta_k. \quad (27)$$

Consequently, we call d_k an h -step, x_k an h -iterate and k , h -iteration.

3.3 Restoration phase

When the iterate is far from feasible region, the f -step may be extremely small or even does not exist, which causes slow convergence or no convergence. To avoid this problem, we set a minimal step length $\alpha_k^{\min}$. If the trial step length α becomes less than $\alpha_k^{\min}$, we terminate the line search and resort to some restoration algorithm. If this happens, the iterate x_k is also called an h -iterate. In our algorithm, the minimal step length is defined as $\alpha_k^{\min} = \min\{\kappa_4, \kappa_5 \theta_k^{\sigma_3}\}$ with $\kappa_4, \kappa_5 \in (0, 1)$ and $\sigma_3 > 1$.

Besides the situation we just mentioned, there are two other cases in which the algorithm turns to a restoration phase. One case is that the subproblem (17) is not consistent, i.e., (17) has no feasible solution. The second case is that problem (17) is very ill-conditioned such that an unusually large step is generated. The step is considered too large if

$$\max\{\|s_k\|, \|t_k\|\} \geq \max\left\{M_\theta, \kappa_6 \frac{1}{\theta_k^{\sigma_4}}\right\} \quad (28)$$

where M_θ is a large constant and $\sigma_4, \kappa_6 > 0$ are fixed. The main purpose of a restoration algorithm is to find a new point x_k^r which has better feasibility. Formally, point x_k^r should satisfy

$$\theta(x_k^r) \leq \kappa_7 \theta_k^{\max}$$

with $\kappa_7 \in (0, 1)$ fixed.

In feasibility restoration phase, we use an elastic-mode methods to drive iterates closer to feasible region, hoping that a better quadratic model exists near the feasible region. The elastic-mode method is based on solving the following elastic problem

$$\begin{aligned} \min \quad & r^T e + (v + w)^T e + (y + z)^T e + \zeta \\ \text{s.t.} \quad & g(x) \geq -r, \quad h(x) = v - w, \\ & G(x) \geq -y, \quad H(x) \geq -z, \\ & Q(x) \leq \zeta, \\ & r, v, w, y, z, \zeta \geq 0, \end{aligned}$$

where e is a vector of all ones with proper scale. Algorithms for nonlinear programming can be used to solve the above problem, for instance, SQP-based or interior-point-based methods. We do not discuss the details since it is not the main topic of this paper. Now, we have mentioned three different sets of iterations:

$$\begin{aligned} \mathcal{K}_f &= \{k \mid x_k \text{ is an } f\text{-iterate}\}, \\ \mathcal{K}_\theta &= \{k \mid x_k \text{ is an } h\text{-iterate}\}, \\ \mathcal{K}_r &= \{k \mid \text{feasibility restoration phase is called at } x_k\}. \end{aligned}$$

Updating maximal infeasibility $\theta_k^{\max}$ is of essential importance in driving global convergence. To drive the algorithm towards feasibility and, ultimately, to optimality, the sequence $\{\theta_k^{\max}\}$ is required to be non-increasing. Only at h -iterates, the maximal infeasibility is changed. If $k \in \mathcal{K}_\theta \setminus \mathcal{K}_r$, then we update $\theta_k^{\max}$ by

$$\theta_{k+1}^{\max} = \max\{\kappa_8 \theta_k^{\max}, \kappa_9 \theta_k + (1 - \kappa_9) \theta_{k+1}\},$$

where κ_8 and κ_9 are constants in interval $(0, 1)$. Otherwise, if $x_k \in \mathcal{K}_r$, then

$$\theta_{k+1}^{\max} = \kappa_7 \theta_k^{\max},$$

where $\kappa_7 \in (0, 1)$. In summary, after achieving a new iterate, the maximal infeasibility is updated by

$$\theta_{k+1}^{\max} = \begin{cases} \theta_k^{\max}, & k \in \mathcal{K}_f; \\ \kappa_7 \theta_k^{\max}, & k \in \mathcal{K}_r; \\ \theta_{k+1}^{\max} = \max\{\kappa_8 \theta_k^{\max}, \kappa_9 \theta_k + (1 - \kappa_9) \theta_{k+1}\}, & \text{otherwise.} \end{cases} \quad (29)$$

3.4 Detailed description of the algorithm

Now we are ready to present the detailed description of the algorithm, which is given in Algorithm 1.

In Algorithm 1, the algorithm stops when $\theta_k + \|t_k(u)\|$ is close to 0. In fact, we have

Theorem 3.2 *Assume that x is an iterate or an accumulation point of $\{x_k\}$ at which $\theta(x) = 0$ and $t_k(u) = 0$. Then x is an S -stationary point for (1).*

Algorithm 1 Main Algorithm

```

1: Initialization An initial point  $x_0$  and positive symmetric matrix  $B_0$  are given. Choose constants  $M_\theta, \kappa_\theta, \kappa_i$ ,
    $i = 1, 2, \dots, 9, \sigma_i, i = 1, \dots, 5$  and  $\rho > 0$ . Choose  $u > 0$  and  $\theta_0^{\max} > 0$ . Set  $k := 0$ .
2:   repeat
3:     Evaluate functions and their gradients. Determine  $u_k^{\min}$ . Compute  $s_k$ .
4:     Set  $to restoration = 0$  and  $getdirection = 0$ ;
5:     repeat ▷ Searching  $u_k$ .
6:       Solve (17) to obtain a solution  $t_k(u)$ .
7:       if (17) is not consistent or the solution becomes too large in the sense of (28) then
8:          $to restoration = 1$ ;
9:       else
10:        if  $t_k(u)$  satisfies (19) and (20), or it satisfies (21) and (22) then
11:          let  $u_k = u, t_k = t_k(u_k), d_k = s_k + t_k$ , and  $getdirection = 1$ .
12:        else if  $u < u_k^{\min}$  then
13:          let  $u_k = u, t_k = t_k(u_k), d_k = s_k + \gamma_k t_k$ , where  $\gamma_k$  is defined by (24).
14:          Let  $getdirection = 1$ .
15:        else
16:          let  $u = u/2$ .
17:        end if
18:      end if
19:    until  $to restoration = 1$  or  $getdirection = 1$ 
20:    if  $getdirection = 1$  then
21:      Set  $\alpha = 1$  and  $newpoint = 0$ .
22:      if  $s_k, t_k$  satisfy (19) then ▷  $f$ -step
23:        repeat
24:          if  $\alpha$  satisfying (25) and (26) then
25:            Let  $\alpha_k = \alpha, x_{k+1} = x_k + \alpha_k d_k$ .
26:            Update  $\theta_k^{\max}$  to  $\theta_{k+1}^{\max}$  using (29).
27:            Set  $k := k + 1$  and  $newpoint = 1$ ;
28:          else if  $\alpha < \alpha_k^{\min}$  then
29:            Set  $to restoration = 1$ ;
30:          else
31:             $\alpha = \alpha/2$ .
32:          end if
33:        until  $newpoint = 1$  or  $to restoration = 1$ 
34:      else ▷  $h$ -step.
35:        repeat
36:          if  $\alpha$  satisfying (27) then
37:            Let  $\alpha_k = \alpha, x_{k+1} = x_k + \alpha_k d_k$ .
38:            Update  $\theta_k^{\max}$  to  $\theta_{k+1}^{\max}$  using (29).
39:            Set  $k := k + 1$  and  $newpoint = 1$ ;
40:          else
41:             $\alpha = \alpha/2$ .
42:          end if
43:        until  $newpoint = 1$  or  $to restoration = 1$ 
44:      end if
45:    end if
46:    if  $to restoration = 1$  then
47:      Let  $\theta_{k+1}^{\max} = \kappa_7 \theta_k^{\max}$ .
48:      Use restoration phase to find some  $x_k^r$ , such that  $\theta(x_k^r) \leq \theta_k^{\max}$ .
49:      Let  $x_{k+1} = x_k^r, \theta_{k+1}^{\max} = \kappa_7 \theta_k^{\max}$ 
50:    end if
51:  until  $\theta_k + \|t_k(u)\| \leq \epsilon$ 

```

Proof We first consider the case that x is an iterate, for example, $x = x_k$. The equation $\theta_k = 0$ implies that x_k is feasible and that $s_k = 0$. Since $t_k(u) = 0$ solves (17), we have by KKT conditions

$$\nabla f(x_k) - \lambda_k \nabla g(x_k) + \mu_k^T \nabla h(x_k) - \nu_k^T \nabla G(x_k) - \xi_k \nabla H(x_k) = 0,$$

with nonnegative Lagrangian multipliers λ_k , ν_k and ξ_k and a Lagrangian multiplier μ_k for equality constraints. Then x_k is an S-stationary point.

Now consider the case where x is an accumulation point of $\{x_k\}$. Let $\{(\lambda_k, \mu_k, \nu_k, \xi_k)\}$, where λ_k , ν_k and ξ_k are nonnegative, be the corresponding Lagrangian multipliers for the constraints in (17). Denote

$$\eta_k = \frac{1}{u_k} \nabla Q_k^T t_k \quad (30)$$

By Lemmas 4.8 and 4.9, which we will introduce later, the set $\{(t_k, \eta_k, \lambda_k, \mu_k, \nu_k, \xi_k)\}$ is bounded. Without loss of generality, we assume that

$$(x_k, t_k, \eta_k, \lambda_k, \mu_k, \nu_k, \xi_k) \rightarrow (x, t, \eta, \lambda, \mu, \nu, \xi), \text{ and } t_k \rightarrow 0, \text{ as } k \rightarrow \infty.$$

Since $t_k = t_k(u_k)$ solves (17), it follows from the first order necessary for problem (17) that

$$\begin{aligned} \nabla f_k + B_k s_k + B_k t_k + \eta_k \nabla Q_k - \nabla g_k \lambda_k \\ + \nabla h_k \mu_k - \nabla G_k \nu_k - \nabla H_k \xi_k = 0. \end{aligned} \quad (31)$$

Using $\nabla Q_k = \nabla G_k H_k + \nabla H_k G_k$, we have

$$\nabla f_k + B_k s_k + B_k t_k - \nabla g_k \lambda_k + \nabla h_k \mu_k - \nabla G_k (\nu_k - \eta_k H_k) - \nabla H_k (\xi_k - \eta_k G_k) = 0. \quad (32)$$

Let $\hat{\nu}_k = \nu_k - \eta_k H_k$, $\hat{\xi}_k = \xi_k - \eta_k G_k$. Taking limit on both side of (31), we have

$$\nabla f(x) - \nabla g(x) \lambda + \nabla h(x) \mu - \nabla G(x) \hat{\nu} - \nabla H(x) \hat{\xi} = 0.$$

For any $i \in I_G \cap I_H$, we have

$$\begin{aligned} \hat{\nu}^{(i)} &= \lim_{k \rightarrow \infty} \hat{\nu}_k^{(i)} = \lim_{k \rightarrow \infty} (\nu_k^{(i)} - \eta_k H_k^{(i)}) = \nu^{(i)} \geq 0, \\ \hat{\xi}^{(i)} &= \lim_{k \rightarrow \infty} \hat{\xi}_k^{(i)} = \lim_{k \rightarrow \infty} (\xi_k^{(i)} - \eta_k G_k^{(i)}) = \xi^{(i)} \geq 0, \end{aligned}$$

which concludes the S-stationarity of x . $\square$

4 Global convergence

For the analysis of global convergence, we make the following assumptions.

Assumptions A:

(A1) There exists a bounded closed convex set $\Omega \subset R^n$ such that $\{x_k\} \subset \Omega$.

(A2) The mappings f , g , h , G , H are continuously differentiable. And the gradients ∇f , ∇g , ∇h , ∇G , ∇H are Lipschitz continuous on Ω .

(A3) The sequenc $\{B_k\}$ is uniformly positive definite and bounded, i.e., there exist positive constants ζ and M such that $d^T B_k d \geq \zeta \|d\|^2$ and $\|B_k\| \leq M$ for all k and d .

Follows from these assumptions, the sequences $\{f_k\}$, $\{g_k\}$, $\{h_k\}$, $\{G_k\}$, $\{H_k\}$, $\{\nabla f_k\}$, $\{\nabla g_k\}$, $\{\nabla h_k\}$, $\{\nabla G_k\}$, $\{\nabla H_k\}$ are bounded. For brevity, we denote with M the upper bound of their ℓ_2 norms. We also use $L > 0$ to denote the Lipschitz constant for all the gradient mappings on Ω .

From the mechanism of inner loops for searching an acceptable step length at an f-iteration, the inner loop will either find an acceptable step length finitely or turns to restoration phase because of too small step length. Now, let us consider the h-iterations. The next lemma relates the estimation of the infeasibility of the trial points at h-iterates.

Lemma 4.1 *Assume that s_k and t_k are directions obtained from (13) and (17), and that (22) holds. Then, for any $\alpha \in (0, 1]$, we have*

$$\theta(x_k + \alpha d_k) \leq (1 - \alpha(1 - 5\kappa_3))\theta_k + \alpha^2 L(\|s_k\| + \|t_k\|)^2. \quad (33)$$

Proof First, we consider θ^C . Using the mean-value theorem and the Lipschitz continuity, we obtain that

$$\begin{aligned} & |\theta^C(x_k + \alpha d_k) - |Q(x_k) + \alpha \nabla Q_k^T d_k|| \\ & \leq \sup_{x_k^\tau \in [x_k, x_k + \alpha d_k]} ||Q_k + \alpha \nabla Q_k(x_k^\tau)^T d_k| - |Q(x_k) + \alpha \nabla Q_k^T d_k|| \\ & \leq \sup_{x_k^\tau \in [x_k, x_k + \alpha d_k]} |Q_k + \alpha \nabla Q_k(x_k^\tau)^T d_k - Q_k - \alpha \nabla Q_k(x_k)^T d_k| \\ & \leq L\|\alpha d_k\|^2 = \alpha^2 L\|s_k + \gamma_k t_k\|^2 \leq \alpha^2 L(\|s_k\| + \|t_k\|)^2. \end{aligned}$$

Using this inequality, the definition of s_k , and (22), we have

$$\begin{aligned} \theta^C(x_k + \alpha d_k) & \leq |Q(x_k) + \alpha \nabla Q_k^T d_k| + \alpha^2 L(\|s_k\| + \|t_k\|)^2 \\ & \leq |Q(x_k) + \alpha \nabla Q_k^T s_k| + |\alpha \gamma_k \nabla Q_k t_k| + \alpha^2 L(\|s_k\| + \|t_k\|)^2 \\ & \leq |Q(x_k) + \alpha \nabla Q_k^T s_k| + \alpha |\nabla Q_k t_k| + \alpha^2 L(\|s_k\| + \|t_k\|)^2 \\ & \leq (1 - \alpha)\theta_k^C + \alpha \kappa_3 \min\{\theta_k^{\sigma_2-1}, \kappa_2\}\theta_k + \alpha^2 L(\|s_k\| + \|t_k\|)^2. \end{aligned}$$

Then, we consider θ^F . Further applications of the mean-value theorem give that

$$\begin{aligned} & ||(g(x_k + \alpha d_k))^- - (g(x_k) + \nabla g_k^T \alpha d_k)^-|| \\ & \leq ||(g(x_k + \alpha d_k) - (g(x_k) + \nabla g_k^T \alpha d_k))^-|| \\ & \leq \sup_{x_k^\tau \in [x_k, x_k + \alpha d_k]} ||(g(x_k) + \nabla g(x_k^\tau)^T \alpha d_k - g(x_k) - \nabla g_k^T \alpha d_k)^-|| \\ & \leq L\|\alpha d_k\|^2 = \alpha^2 L(\|s_k\| + \|t_k\|)^2. \end{aligned}$$

Using this inequality and the constraints of (17), we have

$$\begin{aligned} \|g(x_k + \alpha d_k)^-\| & \leq \|(g(x_k) + \nabla g_k^T \alpha d_k)^-\| + \alpha^2 L(\|s_k\| + \|t_k\|)^2 \\ & \leq \|(g(x_k) + \alpha \nabla g_k^T s_k + \alpha \gamma_k \nabla g_k^T t_k)^-\| + \alpha^2 L(\|s_k\| + \|t_k\|)^2 \\ & \leq \|(g(x_k) + \alpha \nabla g_k^T s_k + \alpha \nabla g_k^T t_k - \alpha(1 - \gamma_k) \nabla g_k^T t_k)^-\| + \alpha^2 L(\|s_k\| + \|t_k\|)^2 \\ & \leq (1 - \alpha)\|(g(x_k))^-\| + \alpha(1 - \gamma_k)\|\nabla g_k^T t_k\| + \alpha^2 L(\|s_k\| + \|t_k\|)^2 \\ & \leq (1 - \alpha)\|g(x_k)^-\| + \alpha(1 - \gamma_k)\kappa_3 \min\{\theta_k^{\sigma_2-1}, \kappa_2\}\theta_k + \alpha^2 L(\|s_k\| + \|t_k\|)^2 \end{aligned}$$

By similar deductions, we obtain that

$$\begin{aligned} \|h(x_k + \alpha d_k)\| & \leq (1 - \alpha)\|h(x_k)\| + \alpha(1 - \gamma_k)\kappa_3 \min\{\theta_k^{\sigma_2-1}, \kappa_2\}\theta_k + \alpha^2 L(\|s_k\| + \|t_k\|)^2, \\ \|G(x_k + \alpha d_k)^-\| & \leq (1 - \alpha)\|G(x_k)^-\| + \alpha(1 - \gamma_k)\kappa_3 \min\{\theta_k^{\sigma_2-1}, \kappa_2\}\theta_k + \alpha^2 L(\|s_k\| + \|t_k\|)^2, \\ \|H(x_k + \alpha d_k)^-\| & \leq (1 - \alpha)\|H(x_k)^-\| + \alpha(1 - \gamma_k)\kappa_3 \min\{\theta_k^{\sigma_2-1}, \kappa_2\}\theta_k + \alpha^2 L(\|s_k\| + \|t_k\|)^2. \end{aligned}$$

Then,

$$\begin{aligned}
\theta(x_k + \alpha d_k) &= \theta^C(x_k + \alpha d_k) + \theta^F(x_k + \alpha d_k) \\
&\leq (1 - \alpha)\theta_k + 5\alpha\kappa_3 \min\{\theta_k^{\sigma_2-1}, \kappa_2\}\theta_k + \alpha^2 L(\|s_k\| + \|t_k\|)^2 \\
&\leq (1 - \alpha)\theta_k + 5\alpha\kappa_3\theta_k + \alpha^2 L(\|s_k\| + \|t_k\|)^2 \\
&\leq (1 - \alpha(1 - 5\kappa_3))\theta_k + \alpha^2 L(\|s_k\| + \|t_k\|)^2.
\end{aligned}$$

□

After similar arguments, we have a similar result in the case where (20) holds.

Lemma 4.2 *Assume that s_k and t_k are directions obtained from (13) and (17) and that (20) holds. Then, for any $\alpha \in (0, 1]$, we obtained that*

$$\theta(x_k + \alpha d_k) \leq (1 - \alpha)\theta_k + 5\alpha\kappa_1\theta_k^{\max} + \alpha^2 L(\|s_k\| + \|t_k\|)^2. \quad (34)$$

The following result is a direct consequence of Lemma 4.1.

Lemma 4.3 *Assume that the subproblem is consistent at x_k , conditions (21) and (22) hold and that*

$$0 < \alpha \leq \alpha_k^h \stackrel{\text{def}}{=} \min \left\{ 1, \frac{(1 - 5\kappa_3 - \rho)\theta_k}{L(\|s_k\| + \|t_k\|)^2} \right\}. \quad (35)$$

Then (27) holds. Hence, the line search at h -iterations can always terminate finitely.

Proof Note that $\theta_k > 0$ at any h -iterate. Using (33) and (35), we obtain that

$$\theta(x_k + \alpha d_k) \leq (1 - \alpha(1 - 5\kappa_3))\theta_k + \alpha^2 L(\|s_k\| + \|t_k\|)^2 \leq (1 - \alpha\rho)\theta_k.$$

□

We recall a result concerning a direct consequence of the definition of $\theta_k^{\max}$ [28].

Lemma 4.4 *The sequence $\{\theta_k^{\max}\}$ is non-increasing and the inequality*

$$0 \leq \theta_l \leq \theta_k^{\max}$$

holds for all $l \geq k$.

The most important role of the maximal infeasibility is to drive the iterates to the feasible region.

Theorem 4.1 *Suppose that $|\mathcal{K}_\theta| = \infty$, where $|\cdot|$ represents the cardinal number of a set. Then $\lim_{k \rightarrow \infty} \theta_k = 0$.*

Proof For the proof, we define

$$\bar{\mathcal{K}}_\theta = \{ k \mid \theta_{k+1}^{\max} = \kappa_9\theta_k + (1 - \kappa_9)\theta_{k+1} \} \setminus \mathcal{K}_r.$$

Assume that $|\mathcal{K}_\theta \setminus \bar{\mathcal{K}}_\theta| = \infty$. By the updating rules of $\theta_k^{\max}$ and Lemma 4.4, we obtain that

$$\begin{aligned}
\theta_{k+1}^{\max} &= \kappa_8\theta_k^{\max} \text{ or } \kappa_7\theta_k^{\max}, \quad k \in \mathcal{K}_\theta \setminus \bar{\mathcal{K}}_\theta, \\
\theta_{k+1}^{\max} &\leq \theta_k^{\max}, \quad k \in \bar{\mathcal{K}}_\theta.
\end{aligned}$$

Then it follows that $\lim_{k \rightarrow \infty} \theta_k^{\max} = 0$, which implies $\lim_{k \rightarrow \infty} \theta_k = 0$.

Now consider the alternative case in which $|\mathcal{K}_\theta \setminus \bar{\mathcal{K}}_\theta| < \infty$. In this case, there exists an index k_0 such that for all $k \in \mathcal{K}_\theta$ and $k \geq k_0$, the feasibility restoration phase is not called and

$$\theta_{k+1}^{\max} = \kappa_9 \theta_k + (1 - \kappa_9) \theta_{k+1}. \quad (36)$$

After simple calculation, we obtain that

$$\theta_k^{\max} - \theta_{k+1}^{\max} \geq \theta_k - \theta_{k+1}^{\max} \geq (1 - \kappa_9) \rho \alpha_k \theta_k.$$

From (28) and (35), it follows that

$$\begin{aligned} \theta_k^{\max} - \theta_{k+1}^{\max} &\geq (1 - \kappa_9) \rho \alpha_k \theta_k \\ &\geq \frac{1}{2} (1 - \kappa_9) \rho \theta_k \min \left\{ 1, \frac{(1 - 5\kappa_3 - \rho) \theta_k}{L(\|s_k\| + \|t_k\|)^2} \right\}. \end{aligned}$$

Then

$$\theta_k^{\max} - \theta_{k+1}^{\max} \geq \frac{1}{2} (1 - \kappa_9) \rho \theta_k \min \left\{ 1, \frac{4(1 - 5\kappa_3 - \rho) \theta_k}{L} \left(\max \left\{ \frac{1}{M_\theta}, \frac{\theta_k^{\sigma_4}}{\kappa_6} \right\} \right)^2 \right\}.$$

Since the sequence of maximal infeasibility $\{\theta_k^{\max}\}$ is non-increasing and thus convergent, the left-hand-side of this inequality converges to 0, which leads to

$$\lim_{k \in \mathcal{K}_\theta} \theta_k = 0. \quad (37)$$

Using (36) and $\theta_{k+1} \leq \theta_k$, we have

$$\theta_{k+1}^{\max} \leq \theta_k$$

for $k \in \mathcal{K}_\theta$ and $k \geq k_0$. This inequality, combined with (37), yields

$$\lim_{k \in \mathcal{K}_\theta} \theta_{k+1}^{\max} = 0.$$

Then we obtain $\lim_{k \rightarrow \infty} \theta_k^{\max} = 0$ because of non-increasingness of $\{\theta_k^{\max}\}$. Consequently, by Lemma 4.4, we have $\lim_{k \rightarrow \infty} \theta_k = 0$. $\square$

Theorem 4.2 *Assume that the algorithm does not terminate in finite steps and that $|\mathcal{K}_\theta| < \infty$. Then $\lim_{k \rightarrow \infty} \theta_k = 0$.*

Proof Let $\{x_k\}$ be the sequence of iterates generated by the algorithm. By the mechanism of line search, we obtain that $\alpha_k \geq 0.5\alpha_k^{\min}$. Since $\sum_{k=1}^{+\infty} (f_k - f_{k+1})$ is bounded, it follows from (26) and the definition of $\alpha_k^{\min}$ that $\lim_{k \rightarrow \infty} \theta_k = 0$. $\square$

Now we consider the behavior of the algorithm near a feasible point. This is important for our global convergence analysis.

Lemma 4.5 *Assume that $\tilde{x}$ is feasible point at which MPEC-MFCQ holds. Then there exists a neighborhood $\tilde{\mathcal{N}}$ of $\tilde{x}$ such that for all $x_k \in \tilde{\mathcal{N}}$, the problem (17) is consistent.*

Proof The main ideas of this proof come from the proof of Lemma 5 in [35] by Fletcher et al. Since $\tilde{x}$ is a feasible point, by MPEC-MFCQ, there exists a vector $\tilde{d}$ such that (8)-(12) hold for $d = \tilde{d}$. Since, by (12), the vectors

$$\begin{aligned} \nabla G^{(i)}(\tilde{x}), \quad i \in I_G \setminus I_H, \quad \nabla H^{(i)}(\tilde{x}), \quad i \in I_H \setminus I_G, \\ \nabla H^{(i)}(\tilde{x}), \quad i = 1, 2, \dots, p \end{aligned}$$

are linearly independent, by continuity, there is a neighborhood $\tilde{\mathcal{N}}$ of $\tilde{x}$, such that for all $x_k \in \tilde{\mathcal{N}}$, the vectors

$$\begin{aligned} \nabla G^{(i)}(x_k), \quad i \in I_G \setminus I_H, \quad \nabla H^{(i)}(x_k), \quad i \in I_H \setminus I_G, \\ \nabla H^{(i)}(x_k), \quad i = 1, 2, \dots, p \end{aligned}$$

are also linearly independent. For brevity, we define a vector-valued functions $C(x)$ which comprises all the constraints involved above. Note that

$$p_k = -(\nabla C_k)^\dagger{}^T (C_k + \nabla C_k^T s_k)$$

satisfies $C_k + \nabla C_k^T(s_k + t) = 0$, where, $(\nabla C_k)^\dagger = (\nabla C_k^T \nabla C_k)^{-1} \nabla C_k^T$, and that

$$\|p_k\| = O(\|s_k\| + \|C_k\|). \quad (38)$$

Let

$$q_k = \frac{(I - \nabla C_k \nabla C_k^\dagger) \tilde{d}}{\|(I - \nabla C_k \nabla C_k^\dagger)\|},$$

where (8) and (9) are used in the second equation. By (8)-(12) and continuity, there is a (smaller) neighborhood $\tilde{\mathcal{N}}$ and a constant $a_1 > 0$ such that

$$\nabla g^{(i)}(x_k)^T q_k \geq a_1, \quad i \in I_g; \quad (39)$$

$$\nabla G^{(i)}(x_k)^T q_k \geq a_1, \quad \nabla H^{(i)}(x_k)^T q_k \geq a_1, \quad i \in I_G \cap I_H; \quad (40)$$

We now consider the solution of (17) and in particular, the line segment defined by

$$v_k(\beta) = p_k + \beta q_k, \quad \beta \in [0, 1]. \quad (41)$$

Clearly, $v_k(\beta)$ satisfies the constraints $C_k + \nabla C_k^T(s_k + t) = 0$. For an active constraint $i \in I_g$, we have by (39) and (41)

$$\begin{aligned} g^{(i)}(x_k) + \nabla g^{(i)}(x_k)^T v_k(\beta) \\ = g^{(i)}(x_k) + \nabla g^{(i)}(x_k)^T p_k + \beta \nabla g^{(i)}(x_k)^T q_k \\ \geq g^{(i)}(x_k) + \nabla g^{(i)}(x_k)^T p_k + \beta a_1. \end{aligned}$$

Similarly, for $i \in I_G \cap I_H$, we have by (40) that

$$\begin{aligned} G^{(i)}(x_k) + \nabla G^{(i)}(x_k)^T v_k(\beta) &\geq G^{(i)}(x_k) + \nabla G^{(i)}(x_k)^T p_k + \beta a_1, \\ H^{(i)}(x_k) + \nabla H^{(i)}(x_k)^T v_k(\beta) &\geq H^{(i)}(x_k) + \nabla H^{(i)}(x_k)^T p_k + \beta a_1. \end{aligned}$$

Then it follows that

$$\left. \begin{aligned} g^{(i)}(x_k) + \nabla g^{(i)}(x_k)^T v_k(\beta) &\geq 0, \quad i \in I_g; \\ G^{(i)}(x_k) + \nabla G^{(i)}(x_k)^T v_k(\beta) &\geq 0 \\ H^{(i)}(x_k) + \nabla H^{(i)}(x_k)^T v_k(\beta) &\geq 0 \end{aligned} \right\}, \quad i \in I_G \cap I_H$$

if

$$\beta \geq \frac{1}{a_1} \max \left\{ \begin{aligned} &|g^{(i)}(x_k) + \nabla g^{(i)}(x_k)^T p_k|, i \in I_g; \\ &|G^{(i)}(x_k) + \nabla G^{(i)}(x_k)^T p_k| \\ &|H^{(i)}(x_k) + \nabla H^{(i)}(x_k)^T p_k| \end{aligned} \right\}, i \in I_G \cap I_H \right\}$$

The definitions of s_k and p_k imply that the lower bound given above is of the order $O(\theta_k + \|C_k\|)$. Thus we can choose a positive constant $a_2 > 0$ such that vector $v_k(\beta)$ satisfies all the active constraints for all $\beta \geq a_2(\theta_k + \|C_k\|)/a_1$.

Next we look at the inactive constraints. By continuity, there are positive constants b_1 and b_2 such that for any inactive constraint indexed by i

$$\begin{aligned} g^{(i)}(x_k) &\geq b_1, \quad \nabla g^{(i)}(x_k)^T q_k \geq -b_2; \\ G^{(i)}(x_k) &\geq b_1, \quad \nabla G^{(i)}(x_k)^T q_k \geq -b_2; \\ H^{(i)}(x_k) &\geq b_1, \quad \nabla H^{(i)}(x_k)^T q_k \geq -b_2. \end{aligned}$$

It follows that

$$\begin{aligned} &g^{(i)}(x_k) + \nabla g^{(i)}(x_k)^T v_k(\beta) \\ &= g^{(i)}(x_k) + \nabla g^{(i)}(x_k)^T p_k + \beta \nabla g^{(i)}(x_k)^T q_k \\ &\geq b_1 - \beta b_2 + \nabla g^{(i)}(x_k)^T p_k, \end{aligned}$$

and that

$$\begin{aligned} G^{(i)}(x_k) + \nabla G^{(i)}(x_k)^T v_k(\beta) &\geq b_1 - \beta b_2 + \nabla G^{(i)}(x_k)^T p_k, \\ H^{(i)}(x_k) + \nabla H^{(i)}(x_k)^T v_k(\beta) &\geq b_1 - \beta b_2 + \nabla H^{(i)}(x_k)^T p_k \end{aligned}$$

for the inactive constraints. By (38), there is a (smaller) neighborhood $\tilde{\mathcal{N}}$ such that

$$\begin{aligned} g^{(i)}(x_k) + \nabla g^{(i)}(x_k)^T v_k(\beta) &\geq \frac{b_1}{2} - \beta b_2, \\ G^{(i)}(x_k) + \nabla G^{(i)}(x_k)^T v_k(\beta) &\geq \frac{b_1}{2} - \beta b_2, \\ H^{(i)}(x_k) + \nabla H^{(i)}(x_k)^T v_k(\beta) &\geq \frac{b_1}{2} - \beta b_2. \end{aligned}$$

Then, for the inactive constraints

$$\begin{aligned} g^{(i)}(x_k) + \nabla g^{(i)}(x_k)^T v_k(\beta) &> 0, \\ G^{(i)}(x_k) + \nabla G^{(i)}(x_k)^T v_k(\beta) &> 0, \\ H^{(i)}(x_k) + \nabla H^{(i)}(x_k)^T v_k(\beta) &> 0 \end{aligned}$$

if $\alpha \leq b_1/(4b_2)$.

To sum up, after possibly shrinking $\tilde{\mathcal{N}}$, vector $v_k(\beta)$ is a feasible solution of (17) if

$$\frac{a_2}{a_1}(\theta_k + \|C_k\|) \leq \beta \leq \frac{b_1}{4b_2}. \quad (42)$$

□

Lemma 4.6 *Assume that MPEC-MFCQ is satisfied at a feasible point $\tilde{x}$. Then there exists a neighborhood $\tilde{\mathcal{N}}$ of $\tilde{x}$ such that the set $\{(\nabla Q_k^T)^\dagger \mid x_k \in \tilde{\mathcal{N}}\}$ is bounded. Hence, the direction set $\{s_k \mid x_k \in \tilde{\mathcal{N}}\}$ is also bounded.*

Proof For the feasible point $\tilde{x}$, we have that

$$\begin{aligned}
\nabla Q(\tilde{x}) &= \nabla(G(\tilde{x})^T H(\tilde{x})) \\
&= \nabla G(\tilde{x}) H(\tilde{x}) + \nabla H(\tilde{x}) G(\tilde{x}) \\
&= \sum_{i \in I_G \setminus I_H} \nabla G^{(i)}(\tilde{x}) H^{(i)}(\tilde{x}) + \sum_{i \in I_H \setminus I_G} \nabla H^{(i)}(\tilde{x}) G^{(i)}(\tilde{x}) \\
&\quad \sum_{i \in I_G \cap I_H} (\nabla G^{(i)}(\tilde{x}) H^{(i)}(\tilde{x}) + \nabla H^{(i)}(\tilde{x}) G^{(i)}(\tilde{x})) \\
&= \sum_{i \in I_G \setminus I_H} \nabla G^{(i)}(\tilde{x}) H^{(i)}(\tilde{x}) + \sum_{i \in I_H \setminus I_G} \nabla H^{(i)}(\tilde{x}) G^{(i)}(\tilde{x}).
\end{aligned}$$

Since $\nabla G^{(i)}(\tilde{x})$, $i \in I_G \setminus I_H$ and $\nabla H^{(i)}(\tilde{x})$, $i \in I_H \setminus I_G$ are linearly independent, we have that $\nabla Q(\tilde{x}) \neq 0$. By continuity, there exists a (smaller) neighborhood $\tilde{\mathcal{N}}$ of $\tilde{x}$ and a positive constant a_3 such that $\nabla Q_k^T \nabla Q_k \geq a_3$ for all $x_k \in \tilde{\mathcal{N}}$. Having this lower bound in mind, it follows from Assumptions A that

$$\|(\nabla Q_k^T)^\dagger\| = \|\nabla Q_k (\nabla Q_k^T \nabla Q_k)^{-1}\| \leq \frac{M}{a_3}$$

for any $x_k \in \tilde{\mathcal{N}}$. Hence, from the definition of s_k , it follows that, for all $x_k \in \tilde{\mathcal{N}}$,

$$\|s_k\| \leq \|(\nabla Q_k^T)^\dagger\| \|Q_k\| \leq \|(\nabla Q_k^T)^\dagger\| \|G_k\| \|H_k\| \leq \frac{M^3}{a_3}. \quad (43)$$

□

The next lemma gives the limitation of $\nabla Q_k^T t_k$ as the parameter u_k approaches to 0.

Lemma 4.7 *Assume that MPEC-MFCQ is satisfied at a feasible point $\tilde{x}$. Let $\{x_k\}_{k \in \mathcal{K}}$ be a subsequence of $\{x_k\}$ that converges to $\tilde{x}$. Suppose that $\lim_{k \in \mathcal{K}} u_k = 0$. Then $\lim_{k \in \mathcal{K}} \|\nabla Q_k^T t_k\| = 0$.*

Proof Let $\beta_k = \frac{a_2}{a_1}(\theta_k + \|C_k\|)$. Then by (42) in the proof of Lemma 4.5, vector $v_k = v_k(\beta_k)$ is a feasible solution for (17). By the optimality of t_k , we have

$$\begin{aligned}
&(\nabla f(x_k) + B_k s_k)^T t_k + \frac{1}{2} t_k^T B_k t_k + \frac{1}{2u_k} \|\nabla Q_k^T t_k\|^2 \\
&\leq (\nabla f(x_k) + B_k s_k)^T v_k + \frac{1}{2} v_k^T B_k v_k + \frac{1}{2u_k} \|\nabla Q_k^T v_k\|^2.
\end{aligned} \quad (44)$$

First we suppose that $\{t_k\}$ is bounded. Then from (44) we have

$$\|\nabla Q_k^T t_k\|^2 \leq 2u_k \left((\nabla f(x_k) + B_k s_k)^T (t_k - v_k) + \frac{1}{2} v_k^T B_k v_k - \frac{1}{2} t_k^T B_k t_k \right) + \|\nabla Q_k^T v_k\|^2.$$

Using this, boundedness and the fact that $\lim_{k \in \mathcal{K}} v_k = 0$, we have $\lim_{k \in \mathcal{K}} \|\nabla Q_k^T t_k\| = 0$.

Now, we consider the other case where $\{t_k\}$ is unbounded. Without loss of generality, we assume that $\lim_{k \in \mathcal{K}} \|t_k\| = +\infty$. Suppose, to arrive at a contradiction, that there is an index $\mathcal{K}' \subset \mathcal{K}$ such that $\lim_{k \in \mathcal{K}'} \|\nabla Q_k^T t_k\| = +\infty$. Then, from (44), Assumptions A, we have, for $k \in \mathcal{K}'$ large enough,

$$\begin{aligned}
\frac{1}{4} \zeta \|t_k\|^2 &\leq \frac{1}{4} \zeta \|t_k\|^2 + \frac{1}{2u_k} (\|\nabla Q_k^T t_k\|^2 - \|\nabla Q_k^T v_k\|^2) \\
&\leq (\nabla f(x_k) + B_k s_k)^T t_k + \frac{1}{2} t_k^T B_k t_k + \frac{1}{2u_k} (\|\nabla Q_k^T t_k\|^2 - \|\nabla Q_k^T v_k\|^2) \\
&\leq (\nabla f(x_k) + B_k s_k)^T v_k + \frac{1}{2} v_k^T B_k v_k.
\end{aligned}$$

Taking limits on these inequalities, we get $+\infty \leq 0$, which is a contradiction.

Summarizing all the arguments above, we conclude the proof. $\square$

Let $(\lambda_k, \mu_k, \nu_k, \xi_k)$ be the Lagrangian multipliers for (17) corresponding to the solution t_k . Next, we discuss the boundedness of the entry set $\{t_k, \lambda_k, \mu_k, \nu_k, \xi_k\}$. We will reveal this fact by proving two lemmas concerning whether or not the sequence $\{u_k\}$ is bounded away from 0. First, we consider the easier case where u_k is bounded away from 0.

Lemma 4.8 *Assume that MPEC-MFCQ holds and that $u_k \geq \bar{u}$ for some positive constant $\bar{u}$. Then the sequence $\{(t_k, \lambda_k, \mu_k, \nu_k, \xi_k)\}$ is bounded.*

Proof Assume, without loss of generality, that $\lim_{k \rightarrow \infty} (x_k, B_k, u_k) = (\tilde{x}, B, \bar{u})$. Since $B_k + \frac{1}{u_k} \nabla Q_k \nabla Q_k^T$ is positive definite, t_k is a solution of (17) if and only if it satisfies the following KKT conditions

$$\begin{aligned} \nabla f_k + B_k s_k + \left(B_k + \frac{1}{u_k} \nabla Q_k \nabla Q_k^T \right) t_k - \nabla g_k \lambda_k \\ + \nabla h_k \mu_k - \nabla G_k \nu_k - \nabla H_k \xi_k = 0, \end{aligned} \quad (45)$$

$$g_k + \nabla g_k^T(s_k + t_k) \geq 0, \quad h_k + \nabla h_k^T(s_k + t_k) = 0, \quad (46)$$

$$G_k + \nabla G_k^T(s_k + t_k) \geq 0, \quad H_k + \nabla H_k^T(s_k + t_k) \geq 0, \quad (47)$$

$$\lambda_k^T (g_k + \nabla g_k^T(s_k + t_k)) = 0, \quad (48)$$

$$\nu_k^T (G_k + \nabla G_k^T(s_k + t_k)) = 0, \quad (49)$$

$$\xi_k^T (H_k + \nabla H_k^T(s_k + t_k)) = 0, \quad (50)$$

$$\lambda_k \geq 0, \quad \nu_k \geq 0, \quad \xi_k \geq 0. \quad (51)$$

Assume that the set $\{\|(t_k, \lambda_k, \mu_k, \nu_k, \xi_k)\|\}$ is unbounded. By taking a subsequence, if necessary, we may assume that $\|(t_k, \lambda_k, \mu_k, \nu_k, \xi_k)\| \rightarrow +\infty$. Without loss of generality, we assume that

$$\|(t_k, \lambda_k, \mu_k, \nu_k, \xi_k)\| \neq 0$$

for all k and that the sequence of vector

$$(\bar{t}_k, \bar{\lambda}_k, \bar{\mu}_k, \bar{\nu}_k, \bar{\xi}_k) := \frac{(t_k, \lambda_k, \mu_k, \nu_k, \xi_k)}{\|(t_k, \lambda_k, \mu_k, \nu_k, \xi_k)\|}$$

converges to some

$$(\bar{t}, \bar{\lambda}, \bar{\mu}, \bar{\nu}, \bar{\xi}) \in R \times R^n \times R^m \times R^p \times R^q \times R^q$$

with

$$\|(\bar{t}, \bar{\lambda}, \bar{\mu}, \bar{\nu}, \bar{\xi})\| = 1 \quad (52)$$

and $\bar{\lambda}, \bar{\nu}, \bar{\xi} \geq 0$. Dividing both sides of (45)-(47) by $\|(t_k, \lambda_k, \mu_k, \nu_k, \xi_k)\|$, dividing both sides of (48)-(50) by $\|(t_k, \lambda_k, \mu_k, \nu_k, \xi_k)\|^2$, and taking the limits as $k \rightarrow \infty$, we get that

$$\begin{aligned} \left(B + \frac{1}{\bar{u}} \nabla Q(\tilde{x}) \nabla Q(\tilde{x})^T \right) \bar{t} - \nabla g(\tilde{x}) \bar{\lambda} + \nabla h(\tilde{x}) \bar{\mu} - \nabla G(\tilde{x}) \bar{\nu} - \nabla H(\tilde{x}) \bar{\xi} = 0, \\ \nabla g(\tilde{x})^T \bar{t} \geq 0, \quad \nabla h(\tilde{x})^T \bar{t} = 0, \\ \nabla G(\tilde{x})^T \bar{t} \geq 0, \quad \nabla H(\tilde{x})^T \bar{t} \geq 0, \\ \bar{\lambda}^T \nabla g(\tilde{x})^T \bar{t} = 0, \\ \bar{\nu}^T \nabla G(\tilde{x})^T \bar{t} = 0, \\ \bar{\xi}^T \nabla H(\tilde{x})^T \bar{t} = 0. \end{aligned} \quad (53)$$

This linear system and the nonnegativity of $\bar{\lambda}$, $\bar{\nu}$ and $\bar{\xi}$ indicate that $\bar{t}$ solves

$$\begin{aligned} \min \quad & \frac{1}{2}t^T \left(B + \frac{1}{\bar{u}} \nabla Q(\tilde{x}) \nabla Q(\tilde{x})^T \right) t, \\ \text{s.t.} \quad & \nabla g(\tilde{x})^T t \geq 0, \quad \nabla h(\tilde{x})^T t = 0, \\ & \nabla G(\tilde{x})^T t \geq 0, \\ & \nabla H(\tilde{x})^T t \geq 0. \end{aligned}$$

Then we have from the positive definiteness of $B + \frac{1}{\bar{u}} \nabla Q \nabla Q^T$ that $\bar{t} = 0$. Hence, it follows from (53) that

$$-\nabla g(\tilde{x})\bar{\lambda} + \nabla h(\tilde{x})\bar{\mu} - \nabla G(\tilde{x})\bar{\nu} - \nabla H(\tilde{x})\bar{\xi} = 0. \quad (54)$$

Now consider the following linear system

$$\begin{aligned} & -\nabla g(\tilde{x})\bar{\lambda} - \sum_{i \in I_G \cap I_H} (\bar{\nu}^{(i)} \nabla G^{(i)}(\tilde{x}) + \bar{\xi}^{(i)} \nabla H^{(i)}(\tilde{x})) \\ & + \nabla h(\tilde{x})\bar{\mu} - \sum_{i \in I_G \setminus I_H} \bar{\nu}^{(i)} \nabla G^{(i)} - \sum_{i \in I_H \setminus I_G} \bar{\xi}^{(i)} \nabla H^{(i)} = 0, \end{aligned} \quad (55)$$

$$\bar{\lambda} \geq 0, \quad (56)$$

$$\bar{\nu}^{(i)} \geq 0, \quad i \in I_G \cap I_H \quad (57)$$

$$\bar{\xi}^{(i)} \geq 0, \quad i \in I_G \cap I_H. \quad (58)$$

$$\sum_{i=1}^m \bar{\lambda}^{(i)} + \sum_{i \in I_G \cap I_H} (\bar{\nu}^{(i)} + \bar{\xi}^{(i)}) \neq 0. \quad (59)$$

Since, by MPEC-MFCQ, the system (8)-(11) has a solution, it follows from the alternative theorem that the system (55)-(59) has no solution. Noting that (54) implies (55), and that (56), (57) and (58) are implied by the nonnegativity of $\bar{\lambda}$, $\bar{\nu}$ and $\bar{\xi}$, there must have

$$\sum_{i=1}^m \bar{\lambda}^{(i)} + \sum_{i \in I_G \cap I_H} (\bar{\nu}^{(i)} + \bar{\xi}^{(i)}) = 0.$$

Then (53) yields

$$\nabla h(\tilde{x})\bar{\mu} - \sum_{i \in I_G \setminus I_H} \bar{\nu}^{(i)} \nabla G^{(i)} - \sum_{i \in I_H \setminus I_G} \bar{\xi}^{(i)} \nabla H^{(i)} = 0.$$

By linear independency (12), we have that

$$\bar{\mu} = 0, \quad \bar{\nu}^{(i)} = 0, \quad i \in I_G \setminus I_H, \quad \bar{\xi}^{(i)} = 0, \quad i \in I_H \setminus I_G.$$

This implies $(\bar{t}, \bar{\lambda}, \bar{\mu}, \bar{\nu}, \bar{\xi}) = 0$, which contradicts with (52). So, the proof is done. $\square$

Lemma 4.9 Assume that MPEC-MFCQ holds and that

$$\lim_{k \rightarrow 0} u_k = 0.$$

Then the sequence $\{(t_k, \lambda_k, \mu_k, \nu_k, \xi_k)\}$ is bounded.

Proof Assume, without loss of generality, that $\lim_{k \rightarrow \infty} (x_k, B_k, u_k) = (\tilde{x}, B, 0)$. Since $B_k + \frac{1}{u_k} \nabla Q_k \nabla Q_k^T$ is positive definite, t_k is a solution of (17) if and only if it satisfies the KKT conditions (45)-(51). Let η_k be a scalar defined by (30). The equation (45) can be reformulated as (32). Assume that the set

$\{\|(\eta_k, t_k, \lambda_k, \mu_k, \nu_k, \xi_k)\|\}$ is unbounded. By taking a subsequence, if necessary, we may assume that $\|(\eta_k, t_k, \lambda_k, \mu_k, \nu_k, \xi_k)\| \rightarrow +\infty$. Without loss of generality, we assume that

$$\|(\eta_k, t_k, \lambda_k, \mu_k, \nu_k, \xi_k)\| \neq 0$$

for all k and that the sequence of vector

$$(\bar{\eta}_k, \bar{t}_k, \bar{\lambda}_k, \bar{\mu}_k, \bar{\nu}_k, \bar{\xi}_k) := \frac{\|(\eta_k, t_k, \lambda_k, \mu_k, \nu_k, \xi_k)\|}{\|(\eta_k, t_k, \lambda_k, \mu_k, \nu_k, \xi_k)\|}$$

converges to some

$$(\bar{\eta}, \bar{t}, \bar{\lambda}, \bar{\mu}, \bar{\nu}, \bar{\xi}) \in R \times R^n \times R^m \times R^p \times R^q \times R^q$$

with

$$\|(\bar{\eta}, \bar{t}, \bar{\lambda}, \bar{\mu}, \bar{\nu}, \bar{\xi})\| = 1$$

and $\bar{\lambda}, \bar{\nu}, \bar{\xi} \geq 0$. Dividing both sides of (32), (46) and (47) by $\|(\eta_k, t_k, \lambda_k, \mu_k, \nu_k, \xi_k)\|$, dividing both sides of (48)-(50) by $\|(\eta_k, t_k, \lambda_k, \mu_k, \nu_k, \xi_k)\|^2$, and taking the limits as $k \rightarrow \infty$, we get that

$$B\bar{t} - \nabla g(\tilde{x})\bar{\lambda} + \nabla h(\tilde{x})\bar{\mu} - \nabla G(\tilde{x})(\bar{\nu} - \bar{\eta}H(\tilde{x})) - \nabla H(\tilde{x})(\bar{\xi} - \bar{\eta}G(\tilde{x})) = 0, \quad (60)$$

$$\nabla g(\tilde{x})^T \bar{t} \geq 0, \quad \nabla h(\tilde{x})^T \bar{t} = 0, \quad (61)$$

$$\nabla G(\tilde{x})^T \bar{t} \geq 0, \quad \nabla H(\tilde{x})^T \bar{t} \geq 0, \quad (62)$$

$$\bar{\lambda}^T \nabla g(\tilde{x})^T \bar{t} = 0, \quad (63)$$

$$\bar{\nu}^T \nabla G(\tilde{x})^T \bar{t} = 0, \quad (64)$$

$$\bar{\xi}^T \nabla H(\tilde{x})^T \bar{t} = 0. \quad (65)$$

Now we need to show that

$$\nabla G^{(i)}(\tilde{x})^T t = 0, \quad i \in I_G \setminus I_H, \quad (66)$$

$$\nabla H^{(i)}(\tilde{x})^T t = 0, \quad i \in I_H \setminus I_G. \quad (67)$$

If this is not true, then at least one of these equalities does not hold. Without loss of generality, let $\nabla G^{(i_0)}(\tilde{x})^T t \neq 0, i_0 \in I_G \setminus I_H$. Then $\nabla G^{(i_0)}(\tilde{x})^T t > 0$ because

$$\nabla G^{(i)}(\tilde{x})^T t = \lim_{k \rightarrow \infty} G^{(i)}(x_k) + \nabla G^{(i)}(x_k)^T t_k \geq 0, \quad \text{for } i \in I_G.$$

Hence, we have

$$\nabla Q(\tilde{x})^T t = \sum_{i \in I_G \setminus I_H} H^{(i)}(\tilde{x}) \nabla G^{(i)}(\tilde{x})^T t + \sum_{i \in I_H \setminus I_G} G^{(i)}(\tilde{x}) \nabla H^{(i)}(\tilde{x})^T t \geq H^{(i_0)}(\tilde{x}) \nabla G^{(i_0)}(\tilde{x})^T t > 0,$$

where the facts that $\nabla H^{(i)}(\tilde{x})^T t \geq 0, i \in I_H$ and $H^{(i)}(\tilde{x}) > 0, i \in I_G \setminus I_H$ are used. This contradicts with Lemma 4.7.

Let

$$\begin{aligned} \hat{\nu} &= \bar{\nu} - \bar{\eta}H(\tilde{x}), \\ \hat{\xi} &= \bar{\xi} - \bar{\eta}G(\tilde{x}). \end{aligned}$$

Clearly, $\hat{\nu}^{(i)} = \bar{\nu}^{(i)}$, $\hat{\xi}^{(i)} = \bar{\xi}^{(i)}$ for all $i \in I_G \cap I_H$. By (66) and (67), we can reformulating system (60)-(65) into

$$\begin{aligned} B\bar{t} - \nabla g(\tilde{x})\bar{\lambda} + \nabla h(\tilde{x})\bar{\mu} - \nabla G(\tilde{x})\bar{\nu} - \nabla H(\tilde{x})\bar{\xi} &= 0, \\ \nabla g(\tilde{x})^T \bar{t} &\geq 0, \quad \nabla h(\tilde{x})^T \bar{t} = 0, \\ \nabla G^{(i)}(\tilde{x})^T \bar{t} &= 0, \quad i \in I_G \setminus I_H, \\ \nabla H^{(i)}(\tilde{x})^T \bar{t} &= 0, \quad i \in I_H \setminus I_G, \\ \nabla G^{(i)}(\tilde{x})^T \bar{t} &= 0, \quad \nabla H^{(i)}(\tilde{x})^T \bar{t} = 0, \quad i \in I_G \cap I_H, \\ \bar{\lambda}^T \nabla g(\tilde{x})^T \bar{t} &= 0, \\ (\hat{\nu}^{(i)}) \nabla G^{(i)}(\tilde{x})^T \bar{t} &= 0, \quad (\hat{\xi}^{(i)}) \nabla H^{(i)}(\tilde{x})^T \bar{t} = 0, \quad i \in I_G \cap I_H. \end{aligned}$$

This system and nonnegativeness of $\hat{\nu}^{(i)}$, $\hat{\xi}^{(i)}$, $i \in I_G \cap I_H$ indicate that $\bar{t}$ solves

$$\begin{aligned} \min \quad & \frac{1}{2} t^T B t, \\ \text{s.t.} \quad & \nabla g(\tilde{x})^T t \geq 0, \quad \nabla h(\tilde{x})^T t = 0, \\ & \nabla G^{(i)}(\tilde{x})^T t = 0, \quad i \in I_G \setminus I_H, \\ & \nabla H^{(i)}(\tilde{x})^T t = 0, \quad i \in I_H \setminus I_G, \\ & \nabla G^{(i)}(\tilde{x})^T t \geq 0, \quad \nabla H^{(i)}(\tilde{x})^T t = 0, \quad i \in I_G \cap I_H. \end{aligned}$$

By the positive definiteness of B , we have $\bar{t} = 0$. Then we get a contradiction after identical discussions as the proof of the previous lemma. $\square$

Besides showing the boundedness of $\{t_k, \lambda_k, \mu_k, \nu_k, \xi_k\}$, Lemmas 4.8 and 4.9 imply the boundedness of $\eta_k = \frac{1}{u_k} \nabla Q_k^T t_k$. This boundedness result guarantees the existence of u_k , with which the corresponding step $t_k(u_k)$ satisfies (18). Assume, without loss of generality that

$$\frac{1}{u_k} |\nabla Q_k^T t_k| \leq M,$$

Then (18) is satisfied for

$$u_k \leq \frac{\delta_k}{M}.$$

By this inequality, (23), (18) and the choice of parameter that $\sigma_2 < \sigma_1$, we have the following result.

Lemma 4.10 *Assume that $\tilde{x}$ is feasible point at which MPEC-MFCQ holds. Then there exists a neighborhood $\tilde{\mathcal{N}}$ of $\tilde{x}$ such that for all $x_k \in \tilde{\mathcal{N}}$, the parameter u_k satisfies $u_k > u_k^{\min}$.*

This lemma ensures that $d_k = s_k + t_k$ near the feasible point. This help us understand the behaviors of the algorithm near points which is feasible but not S-stationary.

Lemma 4.11 *Assume that $\tilde{x}$ is feasible point at which MPEC-MFCQ holds but which is not an S-stationary point. Then there exists a neighborhood $\tilde{\mathcal{N}}$ of $\tilde{x}$ and a positive constant $\bar{\chi}$ such that for all $x_k \in \tilde{\mathcal{N}}$, the solution t_k to (17) satisfies*

$$-\nabla f_k^T t_k > \bar{\chi}. \quad (68)$$

Proof Since t_k solves (17), we have the KKT system (45)-(51). Taking inner product of (47) and t_k , we have

$$\begin{aligned} \nabla f_k^T t_k + s_k^T B_k t_k + t_k^T \left(B_k + \frac{1}{u_k} \nabla Q_k \nabla Q_k^T \right) t_k \\ - \lambda_k^T \nabla g_k^T t_k + \mu_k^T \nabla h_k^T t_k - \nu_k^T \nabla G_k^T t_k - \xi_k^T \nabla H_k^T t_k = 0 \end{aligned}$$

Using (46)-(49) and Assumption (A3), this equality leads to

$$\begin{aligned} -\nabla f_k^T t_k &= t_k^T \left(B_k + \frac{1}{u_k} \nabla Q_k \nabla Q_k^T \right) t_k + s_k^T B_k t_k + \lambda_k^T g_k + \mu_k^T h_k + \nu_k^T G_k + \xi_k^T H_k \\ &\geq \zeta \|t_k\|^2 + s_k^T B_k t_k + \lambda_k^T g_k + \mu_k^T h_k + \nu_k^T G_k + \xi_k^T H_k. \end{aligned} \quad (69)$$

Since $\tilde{x}$ is feasible but not S-stationary, any accumulation point $\tilde{x}$ of $\{t_k\}$ is not 0. Without loss of generality, we assume that

$$(t_k, x_k, \lambda_k, \mu_k, \nu_k, \xi_k) \rightarrow (\tilde{t}, \tilde{x}, \tilde{\lambda}, \tilde{\mu}, \tilde{\nu}, \tilde{\xi}).$$

Then $\tilde{\lambda}, \tilde{\nu}, \tilde{\xi}$ are non-negative vectors because of (51). Using the feasibility of $\tilde{x}$ and continuity, the inequality (69) indicates that there is a (smaller) neighborhood $\tilde{\mathcal{N}}$ such that

$$-\nabla f_k^T t_k \geq \frac{1}{2} \zeta \|\tilde{t}\|^2$$

for all $x_k \in \tilde{\mathcal{N}}$, which concludes the proof with $\bar{\chi} = \zeta \|\tilde{t}\|^2 > 0$. $\square$

Lemma 4.12 *Assume that $\tilde{x}$ is a feasible point at which MPEC-MFCQ hold, and which is not an S-stationary point. Then there exists a (smaller) neighborhood $\tilde{\mathcal{N}}$ of $\tilde{x}$ such that all $x_k \in \tilde{\mathcal{N}}$ satisfy condition (19), and that the step αd_k satisfies condition (25) if α is small enough.*

Proof Choose a neighborhood $\mathcal{N}$ which is contained by the neighborhoods introduced in Lemma 4.10 and 4.11. By Lemma 4.10, we have $d_k = s_k + t_k$. Noting that $|\nabla f(x_k)^T s_k| = O(\theta_k)$, there is a (smaller) neighborhood $\tilde{\mathcal{N}}$ of $\tilde{x}$ such that

$$-\nabla f(x_k)^T d_k = -\nabla f(x_k)^T s_k - \nabla f(x_k)^T t_k \geq \frac{1}{2} \bar{\chi}. \quad (70)$$

Let

$$0 < \alpha \leq \alpha^{f_1} := \frac{(1 - \rho) a_3^2 \bar{\chi}}{4L(M^3 + M a_3)^2}. \quad (71)$$

Using the mean value theorem, Lemma 4.11, Lipschitz continuity, (70), (43) and the upper bound of $\|t_k\|$, we get that

$$\begin{aligned} &f_k - f(x_k + \alpha d_k) - (-\rho \nabla f_k^T \alpha d_k) \\ &\geq -\alpha(1 - \rho) \nabla f(x_k)^T d_k - \sup_{x_k^T \in [x_k, x_k + \alpha d_k]} |(\nabla f(x_k^T) - \alpha \nabla f(x_k))^T d_k| \\ &\geq \frac{1}{2} \alpha(1 - \rho) \bar{\chi} - \alpha^2 L \|d_k\|^2 \\ &\geq \frac{1}{2} \alpha(1 - \rho) \bar{\chi} - \alpha^2 L (\|s_k\| + \|t_k\|)^2 \\ &\geq \frac{1 - \rho}{4} \alpha \bar{\chi}, \end{aligned}$$

which yields

$$f(x_k) - f(x_k + \alpha d_k) \geq -\rho \alpha \nabla f_k^T d_k.$$

$\square$

Lemma 4.13 *Suppose that $\tilde{x}$ is a feasible point at which MPEC-MFCQ holds and which is not an S-stationary point. Then there exists a neighborhood $\tilde{\mathcal{N}}$ of $\tilde{x}$, such that for all $x_k \in \tilde{\mathcal{N}}$, condition (26) is satisfied if*

$$0 < \alpha \leq \alpha_k^{f_2} \triangleq \min \left\{ \frac{(1 - 5\kappa_1) a_3^2}{2L(M^3 + a_3 M)^2} \theta_k^{\max}, \sqrt{\frac{a_3 \theta_k^{\max}}{2L(M^3 + a_3 M)^2}} \right\} \quad (72)$$

Proof First, suppose that $\theta_k \geq \frac{1+5\kappa_1}{2}\theta_k^{\max}$. By Lemma 34, we have

$$\begin{aligned}\theta(x_k + \alpha d_k) &\leq (1 - \alpha)\theta_k + 5\alpha\kappa_1\theta_k^{\max} + \alpha^2 L(\|s_k\| + \|t_k\|)^2 \\ &\leq \theta_k - \alpha(\theta_k - 5\kappa_1\theta_k^{\max}) + \alpha^2 L(\|s_k\| + \|t_k\|)^2 \\ &\leq \theta_k - \alpha \left(\frac{1+5\kappa_1}{2}\theta_k^{\max} - 5\kappa_1\theta_k^{\max} \right) + \alpha^2 L(\|s_k\| + \|t_k\|)^2 \\ &\leq \theta_k - \alpha \left(\frac{1-5\kappa_1}{2}\theta_k^{\max} \right) + \alpha^2 L \left(\frac{M^3}{a_3} + M \right)^2,\end{aligned}$$

which yields that

$$\theta(x_k + \alpha d_k) \leq \theta_k \leq \theta_k^{\max}$$

whenever

$$0 < \alpha \leq \frac{(1-5\kappa_1)a_3^2}{2L(M^3 + a_3M)^2}\theta_k^{\max}.$$

Now, consider the other case in which $\theta_k < \frac{1+5\kappa_1}{2}\theta_k^{\max}$. By Lemma 34, we have

$$\begin{aligned}\theta(x_k + \alpha d_k) &\leq (1 - \alpha)\theta_k + 5\alpha\kappa_1\theta_k^{\max} + \alpha^2 L(\|s_k\| + \|t_k\|)^2 \\ &\leq (1 - \alpha)\frac{1+5\kappa_1}{2}\theta_k^{\max} + 5\alpha\kappa_1\theta_k^{\max} + \alpha^2 L(\|s_k\| + \|t_k\|)^2 \\ &\leq \frac{1 - \alpha(1-5\kappa_1)}{2}\theta_k^{\max} + \alpha^2 L(\|s_k\| + \|t_k\|)^2 \\ &\leq \frac{1}{2}\theta_k^{\max} + \alpha^2 L \left(\frac{M^3}{a_3} + M \right)^2.\end{aligned}$$

Hence, condition (26) holds for the step length α satisfying

$$0 < \alpha \leq \sqrt{\frac{a_3\theta_k^{\max}}{2L(M^3 + a_3M)^2}}.$$

To sum up, condition (26) is satisfied for all α satisfying (72) □

Lemma 4.14 *Suppose that $\tilde{x}$ is a feasible point at which MPEC-MFCQ holds and which is not an S -stationary point. Then there exists a smaller neighborhood $\tilde{\mathcal{N}}$ of $\tilde{x}$, such that for all $x_k \in \tilde{\mathcal{N}}$, the corresponding line search will terminate with an acceptable step length, in other words, the algorithm will not turn to restoration phase.*

Proof Firstly, by Lemmas 4.6, 4.8 and 4.9, the direction sets $\{s_k\}$ and $\{t_k\}$ are bounded near the feasible region. Then the switching condition (28) to feasibility restoration will not be activated when x_k is almost feasible.

Secondly, by Lemma 4.5, the subproblem (17) is consistent near $\tilde{x}$, which ensures not calling restoration phase because of inconsistency.

Finally, let us show that feasibility restoration phase will not happen during line search at f -iterate near $\tilde{x}$. Using Lemma 4.12 and 4.13, the line search will find an acceptable step length when $0 < \alpha \leq \min\{\alpha^{f_1}, \alpha_k^{f_2}\}$. By the expression of α^{f_1} and $\alpha_k^{f_2}$, we obtain that $\min\{\alpha^{f_1}, \alpha_k^{f_2}\} \geq O(\theta_k)$. On the other hand, the definition of $\alpha_k^{\min}$ shows $\alpha_k^{\min} = O(\theta_k^{\sigma_3})$. Since $\sigma_3 > 1$, there exists a (smaller) neighborhood $\tilde{\mathcal{N}}$ of $\tilde{x}$ such that for all $x_k \in \tilde{\mathcal{N}}$, there must be $\alpha_k^{\min} < 0.5 \min\{\alpha^{f_1}, \alpha_k^{f_2}\}$, which implies that the Amijio line search procedure finished successfully with the step length $\alpha_k \geq \alpha_k^{\min}$. This also implies that the restoration phase is not called. □

Theorem 4.3 *Assume that $\{x_k\}$ is an infinite sequence generated by the algorithm, $|\mathcal{K}_\theta| = \infty$ and MPEC-MFCQ holds at any feasible points. Then any accumulation point of $\{x_k\}_{k \in \mathcal{K}_\theta}$ is S-stationary.*

Proof We have shown in Theorem 4.1 that $\lim_{k \rightarrow \infty} \theta_k = 0$. Assume, to arrive at a contradiction, that there is an accumulation point $\tilde{x}$ which is not S-stationary. Then, by Lemma (4.11), there is an index set $\mathcal{K}$ and a constant $\bar{\chi} > 0$ such that for sufficiently large $k \in \mathcal{K}$, the inequality (68) holds. On the other hand, from Lemma 4.14, the restoration phase does not happen when $k \in \mathcal{K}$ is sufficiently large. Thus, for sufficiently large $k \in \mathcal{K}_\theta$, it follows the way of updating $\theta_k^{\max}$ that

$$-\nabla f(x_k)^T(s_k + t_k) \leq \kappa_\theta \theta_k,$$

which results in

$$\bar{\chi} \leq -\nabla f(x_k)^T t_k \leq \kappa_\theta \theta_k,$$

where (68) is used. Taking limits on both sides of this inequality, we get that $\bar{\chi} \leq 0$, which is a contradiction. Hence, the proof is done. $\square$

Theorem 4.4 *Assume that $\{x_k\}$ is an infinite sequence generated by the algorithm, $|\mathcal{K}_\theta| < \infty$ and MPEC-MFCQ hold at any feasible points. Then any accumulation point of $\{x_k\}$ is an S-stationary point.*

Proof Assume, for the purpose of deriving a contradiction, that there exists some index set $\mathcal{K}$ such that $\lim_{k \in \mathcal{K}} x_k = \tilde{x}$ and $\tilde{x}$ is not an S-stationary point. Then $\tilde{x}$ is a feasible point by Theorem 4.2. It follows from Lemma 4.5 that

$$-\nabla f_k^T t_k \geq \bar{\chi}$$

for some positive scalar $\bar{\chi}$. Since $|\mathcal{K}_\theta| < +\infty$, there is an index $k_1 > 0$ such that for all $k \geq k_1$, the maximal infeasibility remains unchanged, i.e., $h_k^{\max} = h_{\bar{k}}^{\max}$. If we restrict our attention to $k \in \mathcal{K}$, by Lemma 4.12 and 4.13 and the law of line search, for any $k \geq k_1$, increase k_1 if necessary,

$$\alpha_k \geq 0.5 \min\{\alpha^{f_1}, \alpha_{\bar{k}}^{f_2}\}. \quad (73)$$

It follows, using (71), (72) and (73), from continuity that

$$\begin{aligned} f_k - f_{k+1} &\geq -\alpha_k \rho \nabla f_k^T d_k \\ &= -\alpha_k \rho (\nabla f_k^T s_k + \nabla f_k^T t_k) \\ &\geq \frac{1}{2} \alpha_k \rho \bar{\chi} \\ &\geq \frac{1}{4} \rho \min\{\alpha^{f_1}, \alpha_{\bar{k}}^{f_2}\} \bar{\chi} \end{aligned}$$

Note that the right-hand side of the last inequality is a constant in this case. It follows that

$$\begin{aligned} \sum_{k=0}^{\infty} (f_k - f_{k+1}) &\geq \sum_{k \in \mathcal{K} \text{ and } k \geq k_1} (f_k - f_{k+1}) \\ &\geq \sum_{k \in \mathcal{K} \text{ and } k \geq k_1} \frac{1}{4} \min\{\alpha^{f_1}, \alpha_{\bar{k}}^{f_2}\} \bar{\chi}, \end{aligned}$$

which is a contradiction because the series in the left-hand side of the first inequality is convergent while that in the right-hand side of the second inequality is diverge. $\square$

5 Final remark

In this paper, we propose a new SQP-based framework for solving MPECs. The key point is to view MPEC as combination of two problems which are easier to solve. This idea can be considered as a generalization of the Byrd-Omojokun's trust region strategy. A trust-funnel-like strategy is used to balance the improvements on feasibility and optimality. The presented algorithm is shown to be globally convergent to S-stationary points under MPEC-MFCQ. The global convergence results are satisfactory. Further work will focus on providing local convergence analysis and numerical experiments.

On the other hand, we think that some of the ideas of this algorithm may be useful for designing algorithm for nonlinear programs. For example, we have applied the ideas of computing nonlinear programming step into trust region methods and interior points methods.

References

1. Luo, Z.S., Pang, J.S., Ralph, D.: Mathematical Programs with equilibrium Constraints. Cambridge University Press (1996)
2. Outrata, J., Kočvara, M., Zowe, J.: Nonsmooth Approach to Optimization Problem with Equilibrium Constraints: Theory, Applications and Numerical Results. Kluwer, Dordrecht, The Netherlands (1998)
3. Chen, Y., Flrian, F.: The nonlinear bilevel programming problem: formulations, regularity and optimality conditons. *Optimization* **32**, 193–209 (1995)
4. Ye, J.J., Zhu, D.L., Zhu, Q.L.: Exact penalization and necessary optimality conditions for generalized bilevel programming problems. *SIAM Journal on Optimization* **7**(2), 481–507 (1997).
5. Facchinei, F., Jiang, H.Y., Qi, L.Q.: A smoothing method for mathematical programs with equilibrium constraints. *Mathematical Programming* **85**, 81–106 (1999)
6. Fukushima, M., Pang, J.S.: convergence of a smoothing continuation method for mathematical programs with complementarity constraints. In: M. Thëra, R. Tichatschke (eds.) *Ill-posed variational problems and regularity techniques., Lecture notes in economics and mathematical systems*, vol. 447. Springer, Heidelberg (1999)
7. Scholtes, S.: Convergence properties of a regularization scheme for mathematical programs with complementarity constraints. *SIAM Journal on Optimization* **11**(4), 918–936 (2001).
8. Ralph, D., Wright, S.J.: Some properties of regularization and penalization schemes for mpecs. *Optimization Methods and Software* **19**(5), 527–556 (2004)
9. Hu, X.M., Ralph, D.: Convergence of a penalty method for mathematical programming with complementarity constraints. *J. Optim. Theory Appl.* **123**, 365–369 (2004)
10. Huang, X.X., Yang, X.Q., Zhu, D.L.: A sequential smooth penalization approach to mathematical programs with complementarity constraints. *Numer. Funct. Anal. Optim.* **27**, 71–98 (2006)
11. Jiang, H.Y., Ralph, D.: Extension of quasi-newton methods to mathematical programs with complementarity constraints. *Comput. Optim. Appl.* **25**, 123–150 (2002)
12. Luo, Z.Q., Pang, J.S., Ralph, D.: Piecewise sequential quadratic programming for mathematical programs with nonlinear complementarity constraints. In: A. Migdalas, P. Pardalos, P. Värbrand (eds.) *Multilevel Optimization: Algorithms, Complexity, and Applications*, pp. 209–229. Kluwer Academic Publishers (1998)
13. Scholtes, S., Stöhr, M.: Exact penalization of mathematical programs with complementarity constraints. *SIAM J. Control Optim.* **37**, 617–652 (1999)
14. Anitescu, M.: On using the elastic mode in nonlinear programming approacher to mathematical programs with complementarity constraints. *SIAM J. Optim.* **15**, 1203–1236 (2005)
15. Anitescu, M., Tseng, P., Wright, S.J.: Elastic-mode algorithms for mathematical programs with equilibrium constraints: global convergence and stationarity properties. *Mathematical programming* **110**(2), 337–371 (2007)
16. Fletcher, R., Leyffer, S., Ralph, D., Scholtes, S.: Local convergence of SQP method for mathematical programs with equilibrium constraints. *SIAM J. Optim.* **17**, 259–286 (2006)
17. Liu, X.W., Perakis, G., Sun, J.: A robust SQP method for mathematical programs with linear complementarity constraints. *Comput. Optim. Appl.* **34**, 5–33 (2006)
18. Benson, H.Y., Sen, A., Shanno, D.F., Vanderbei, R.: Interior point algorithms, penalty methods and equilibrium problems. *Comput. Optim. Appl.* **34**, 155–182 (2006)

19. Leyffer, S., López-Calva, G., Nocedal, J.: Interior methods for mathematical programs with complementarity constraints. *SIAM J. Optim.* **17**, 52–77 (2006)
20. Raghunathan, A., Biegler, L.: Interior point methods for mathematical programs with complementarity constraints. *SIAM J. Optim.* **15**, 720–750 (2005)
21. Leyffer, S.: Macmpec: Ampl collections of mpecs, (2005). URL <http://www.msc.anl.gov/~leyffer/MacMPEC>
22. Fletcher, R., Leyffer, S.: Solving mathematical programs with complementarity constraints as nonlinear programs. *Optim. Methods Softw.* **19**, 15–40 (2004)
23. Scheel, H., Scholtes, S.: Mathematical programs with complementarity constraints: Stationarity, optimality, and sensitivity. *Mathematics of Operations Research* **25**(1), 1–22 (2000)
24. Scheel, H., Scholtes, S.: Mathematical programs with complementarity constraints: stationarity, optimality, and sensitivity. *Math. Oper. Res.* **25**, 1–22 (2000)
25. Byrd, R.H., Schnabel, R.B., Shultz, G.A.: A trust region algorithm for nonlinearly constrained optimization. *SIAM J. Numer. Anal.* **24**(5), 1152–1170 (1987).
26. Omojokun, E.: Trust region algorithm for optimization with equalities and inequalities constraints. Ph.D. thesis, Ph. D Thesis, University of Colorado at Boulder (1989)
27. Wright, S.J., Nocedal, J.: Numerical optimization, vol. 2. Springer New York (1999)
28. Gould, N.I.M., Toint, Ph.L.: Nonlinear programming without a penalty function or a filter. *Math. Program.* **122**(1), 155–196 (2010)
29. Bielschowsky, R.H., Gomes, F.A.M.: Dynamic control of infeasibility in equality constrained optimization. *SIAM J Optimization* **19**(3), 1299 – 1325 (2008)
30. Liu, X.W., Yuan, Y.X.: A sequential quadratic programming method without a penalty function or a filter for nonlinear equality constrained optimization. *SIAM J. Optim.* **21**(2), 545–571 (2011)
31. Qiu, S.Q., Chen, Z.W.: Global and local convergence of a class of penalty-free-type methods for nonlinear programming. *Appl. Math. Model.* **36**(7), 3201–3216 (2012)
32. Qiu, S.Q., Chen, Z.W.: A new penalty-free-type algorithm based on trust region techniques. *Appl. Math. Comput.* **218**(22), 11,089–11,099 (2012)
33. Xue, W.J., Shen, C.G., Pu, D.G.: A penalty-function-free line search SQP method for nonlinear programming. *Journal of Computational and Applied Mathematics* **228**(1), 313–325 (2009)
34. Zoppke-Donaldson, C.: A tolerance-tube approach to sequential quadratic programming with applications. Ph.D. thesis, Department of Mathematics and Computer Science, University of Dundee, Dundee, Scotland, UK, (1995)
35. Fletcher, R., Leyffer, S., Toint, P.: On the global convergence of a filter–sqp algorithm. *SIAM J. Optim* **13**(1), 44–59 (2002)